

Supplementary Table S1: A: Dosage table with values in ml for patients not taking stiripentol (from the Fintepla® prescribing information, as of December 2020). B: Dosage table with values in ml for patients taking stiripentol.

A: Recommended titration scheme without stiripentol								
	Dose for 2 x daily administration (mg/kg)	Fenfluramine 2.2 mg/ml = Fenfluramine hydrochloride 2.5 mg/ml						
Weight (kg)		10	15	20	25	30	35	40
Start dose	0.1 mg/kg bw	0.5 ml	0.6 ml	0.8 ml	1.0 ml	1.3 ml	1.5 ml	1.7 ml
Day 7	0.2 mg/kg bw	0.8 ml	1.2 ml	1.6 ml	2.0 ml	2.4 ml	2.8 ml	3.2 ml
Day 14	0.35 mg/kg bw	1.6 ml	2.4 ml	3.2 ml	4.0 ml	4.8 ml	5.6 ml	6.0 ml
	For body weight of 45 kg and above titration can be approximated							
Start dose	Approximately 0.1 mg/kg bw	2 ml						
Day 7	Approximately 0.2 mg/kg bw	4 ml						
Day 14	Approximately 0.35 mg/kg bw	6 ml						

B: Recommended titration scheme with stiripentol

	Dose for 2 x daily administration (mg/kg)	Fenfluramine 2.2 mg/ml = Fenfluramine hydrochloride 2.5 mg/ml						
Weight (kg)		10	15	20	25	30	35	40
Start dose	0.1 mg/kg bw	0.5 ml	0.6 ml	0.8 ml	1.0 ml	1.3 ml	1.5 ml	1.7 ml
Day 7	0.2 mg/kg bw	1.0 ml	1.5 ml	2.0 ml	2.5 ml	3.0 ml	3.6 ml	4.0 ml
	For body weight of 45 kg and above titration can be approximated							
Start dose	Approximately 0.1 mg/kg bw	2 ml						
Day 7	Approximately 0.2 mg/kg bw	4 ml						

*Supplementary Table S2: Reported adverse events during the FFA study (Medical Dictionary for Regulatory Activities, MedDRA) by frequency. MR = Mitral valve regurgitation, AR = Aortic valve regurgitation.*

MedDRA-organ class	Very common	Common
Infections and Parasitic diseases	Bronchitis Upper respiratory tract infection	Ear infection
Metabolic disorders / Feeding disorders	Decreased appetite	
Psychiatric disorders		Abnormal behaviour Irritability
Nervous system disorders	Lethargy Somnolence Status epilepticus	
Gastrointestinal disorders	Obstipation Diarrhoea Vomiting	
General disorders / Discomfort	Pyrexia Fatigue	
Investigations	Lower blood glucose level Abnormal echocardiographic signs (trace regurgitation, MR or AR) Decrease in body weight	
Injury, Intoxication, Procedure-associated complications	Falls	